

Imtrex[®]

Fungicide

Q Why should you use Imtrex[®]?

- A
- Solo SDHI Group 7 fungicide with best-in-class activity on key diseases (e.g. Scald or Speckled leaf blotch)
 - Allows greater choice of MOA partner i.e. can use strobilurin

Q What diseases are controlled by Imtrex?

- A
- | | |
|---|---|
| On Barley: | On Wheat: |
| <ul style="list-style-type: none">■ Scald■ <i>Ramularia</i> (suppression)■ Leaf rust■ Net blotch (spot & net forms)■ Powdery mildew | <ul style="list-style-type: none">■ Speckled leaf blotch (Septoria leaf blotch)■ Brown rust■ Stripe rust■ Powdery mildew |

Q What is Imtrex compatible with?

- A
- Always apply Imtrex in a mixture with another approved, non-cross resistant fungicide recommended for the control of the same target diseases. Refer to manufacturers' recommendations for rates of partner fungicides.
- Can be tank mixed with most commonly used insecticides, herbicides (e.g. Starane[®] or mecoprop-P) and growth regulators (e.g. Cycocel[®] or Terpal[®]).

Q Resistance management?

- A
- Imtrex is a Group 7 (SDHI) fungicide. Do not apply more than two Imtrex fungicide applications per season. For resistance purposes Systiva[®] seed treatment counts as one SDHI fungicide application.

Q What is the withholding period?

- A
- Cereal green feed/silage:** Allow 28 days between last application and introduction of stock for grazing or cutting for silage.
- Cereal grain:** DO NOT harvest within 42 days of last application.
- When mixing observe the longest WHP of any tank mix.**

Imtrex

Fungicide

At a glance

Crops registered in:
Wheat and Barley.

Product type:
Systemic cereal fungicide from the SDHI mode of action chemistry.

Active ingredient:
62.5 g/L fluxapyroxad (SDHI) emulsifiable concentrate.

Pack size:
10 L.

Adjuvant:
Not required.

Rainfastness:
Do not apply if rainfall is likely before Imtrex dries on the crop.

Water rate:
Minimum of 150 L/ha by ground.

Application rate:
Imtrex 1.25 L/ha with 0.625 L/ha Opus.

- Mixing:**
- Add required amount of Imtrex directly into partly filled spray tank with agitation operating. Complete filling and maintain agitation until spraying complete.
 - **DO NOT** leave spray mix standing overnight.

For more information on Imtrex fungicide, visit crop-solutions.basf.co.nz or call 0800 932 273

ALWAYS READ AND FOLLOW LABEL DIRECTIONS BEFORE USING ANY PRODUCT IN THIS FACT SHEET.

Imtrex[®], Opus[®], Cycocel[®], Systiva[®] and Terpal[®] are registered trademarks of BASF. © Copyright BASF 2018 ® Registered trademark of BASF * Registered Trademark.