

WARNING
KEEP OUT OF REACH OF CHILDREN
READ LABEL BEFORE USE

Poncho[®]

Contains 600 g/litre clothianidin in the form of a suspension concentrate

GROUP 4A INSECTICIDE

A systemic insecticide for use as a seed treatment in maize, sweetcorn,
cereal, forage brassica and grass seed crops

Contents: 600 mL, 5 L, 200 L, 1000 L

Registered to and distributed by:

BASF New Zealand Limited

NZBN 9429032151299

P.O. Box 407, Auckland 1140

Phone: 0800 932 273

crop-solutions.basf.co.nz

Hazard Classification

6.1D, 6.3B, 6.4A, 6.5B, 6.9B, 9.1A, 9.2B, 9.3B, 9.4A



WARNING

Harmful if swallowed, in contact with skin, or inhaled. May cause mild skin irritation. May cause serious eye irritation. May cause an allergic skin reaction. May cause respiratory irritation, allergy, asthma symptoms or breathing difficulties if inhaled. May cause organ damage from repeated oral exposure at high doses. Very toxic to aquatic organisms with long-lasting effects. Toxic to the soil environment. Toxic to terrestrial vertebrates. Very toxic to terrestrial invertebrates.

PRECAUTIONS

HANDLING: This product must be under the control of a qualified person during use. Obtain special instructions before use. DO NOT handle until all safety precautions have been read and understood. Use personal protective equipment as required.

STORAGE: Keep out of reach of children. Store in original container tightly closed and in a locked dry, cool, well ventilated area, away from food stuffs.

SPILLAGE: Deal with all spillages immediately. Wear suitable protective clothing as described in the GENERAL SAFETY section. Prevent further spillage and product from entering waterways. Contain the spill by damming, recover spilt product by absorbing with sawdust or an inert absorbent material, then place material in sealable container labelled for disposal. Wash affected area with water and detergent. Absorb and collect washings and place in the same sealable container as the spilled material. Dispose of the waste safely in suitable landfill. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

GENERAL SAFETY AND HYGIENE MEASURES: Do not eat, drink or smoke when using this product. Wear protective clothing such as respiratory protection, impervious gloves, waterproof hat, coat and trousers when using. Remove protective clothing and wash hands and face thoroughly with soap and water before meals and after work. Contaminated work clothing should not be allowed out of the workplace. Wash protective clothing immediately after use. Store work clothing separately.

USE AND ENVIRONMENTAL PROTECTION: Apply with well-maintained and accurately calibrated equipment. DO NOT spray over, or allow drift over, surface water. Clean out application equipment thoroughly after use. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. DO NOT contaminate water when disposing of equipment wash waters.

NOTE: Storage, application and record keeping must be as described in the current version of the New Zealand Standard for the Management of Agrichemicals NZS8409.

DISPOSAL: Product: Dispose of this product only by using according to the label or at an approved landfill. DO NOT burn product. DO NOT contaminate water with product or its container. DO NOT clean application equipment near surface water. Avoid contamination via drains from farmyards and roads. Container: Triple rinse empty container and add rinsate to the seed treatment tank. Recycle through Agrecovery (0800 247 326, www.agrecovery.co.nz).

FIRST AID

If exposed or if you feel unwell, call National Poisons Centre 0800 POISON (0800 764 766) or a doctor/physician. If medical advice is needed, have product container or label at hand. IF SWALLOWED: rinse mouth. DO NOT induce vomiting. IF ON SKIN: remove contaminated clothing and wash with plenty of soap and water. If skin irritation occurs, get medical advice. IF IN EYES: rinse cautiously with water for several minutes. If irritation persists, get medical advice. IF INHALED: remove to fresh air and keep at rest in a position comfortable for breathing. If experiencing respiratory symptoms, get medical advice.

CONDITIONS OF SUPPLY

Read conditions of supply before buying and using. If these conditions are unacceptable, return at once unopened. All goods supplied by BASF New Zealand Limited are of a high grade and conform to the chemical description of the label. However, to the extent permitted by law, neither BASF nor any of its Distributors accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with the supply of those goods including without limitation their storage, application, handling or use. BASF New Zealand Limited nor any of its Distributors, make no warranties of any kind, express or implied, with respect to fitness for purpose (including crop injury, ineffectiveness or any other unintended consequence), merchantable quality or any other matter with respect to these goods. When you are a consumer as defined in the Consumer Guarantees Act 1993 (the "Act") these conditions will be subject to the provisions of the Act. However, the Act will not apply to goods supplied to you for business purposes.

Registered pursuant to the ACVM Act 1997, No. P5967

See www.foodsafety.govt.nz for registration conditions

Approved pursuant to the HSNO Act 1996, Code HSR100825

See www.epa.govt.nz for approval controls

Batch No:

Date of Manufacture:

Registered to and distributed by:

BASF New Zealand Limited

NZBN 9429032151299

P.O. Box 407, Auckland 1140

Phone: 0800 932 273

crop-solutions.basf.co.nz

EMERGENCY PHONELINK
24 Hour Advice in an Emergency Only
0800 944 955



IIIIII BAR CODE IIIIII

GENERAL INFORMATION

Poncho is a systemic insecticidal seed treatment for protection from Argentine stem weevil, black beetle and greasy cutworm in maize and sweetcorn, Argentine stem weevil and grass grub larvae in wheat, aphids in cereals, Argentine stem weevil, springtails and wheat bug in forage brassicas, and Argentine stem weevil, black beetle and grass grub larvae in grass seed.

RESISTANCE MANAGEMENT

Poncho contains clothianidin from the mode of action **GROUP 4A INSECTICIDE**. Repeated use of insecticides with similar modes of action can lead to the build-up of resistant pest populations. The resistant individuals can eventually dominate the insect population. To minimize this risk, read product label before applying any insecticide and follow label instructions. The effectiveness of Poncho on resistant individuals could be significantly reduced. Since occurrence of resistant insects is difficult to detect prior to use, BASF New Zealand Ltd accepts no liability for any losses that may result from the failure of Poncho to control resistant insects. For further information contact your local supplier or a BASF New Zealand Ltd representative. Refer to the NZCPR web site < www.resistance.nzpps.org > for more information.

DIRECTIONS FOR USE

It is an offence to use this product on animals.

Crop	Pest	Rate	Remarks
Maize Sweetcorn	Argentine stem weevil Black beetle Greasy cutworm	100 mL/80,000 seeds.	Mix with sufficient water (1-2 litres/100 kg seed) and apply to seed ensuring even coverage.
Cereals	Aphids	60 mL/100 kg of seed.	Mix with sufficient water (usually 10 litres/tonne of seed) and apply to seed ensuring even coverage.
Wheat	Argentine stem weevil Grass grub larvae	60 mL/100 kg of seed.	Mix with sufficient water (usually 10 litres/tonne of seed) and apply to seed ensuring even coverage.
Forage Brassicas	Argentine stem weevil Springtails (<i>Sminthuridae</i> family) Wheat bug (<i>Nysius huttoni</i>)	12 mL/kg of seed.	Mix with sufficient water (usually 1 litre/100 kg) and apply to seed ensuring even coverage.
Grass seed	Argentine stem weevil Black beetle Grass grub larvae	450 mL/100 kg of seed.	Mix with sufficient water (usually 1 litre/100 kg) and apply to seed ensuring even coverage.

Poncho should only be applied through a properly-calibrated seed treatment machine to high quality undamaged seed. As Poncho treatment may alter the seed flow rate, recalibrate the planter prior to drilling with treated seed to ensure the required plant population is achieved.

COMPATIBILITY:

Poncho can be applied to seed previously treated with a fungicide or in conjunction with a seed treatment fungicide.

WITHHOLDING PERIOD:

Maize, Sweetcorn, Wheat, Cereals, Forage Brassicas and Grass seed	42 days from sowing
Grass seed oversown into pasture (stitching)	21 days from sowing

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

TREATED SEED:

The following statements must be in bold print on the labels of treated seed containers:

- **Treated seed must not be used for animal or human consumption**
- **Treated seed may be harmful to seed-eating birds**
- **Treated seed should be properly sown and completely covered by soil**
- **Excess seed must not be left in areas accessible to birds**
- **Personal protective equipment must be used when handling treated seeds**
- **Unwanted treated seed must be disposed of in accordance with the Hazardous Substances (Disposal) Notice 2017**
- **Withholding period:**
Stock must not graze maize, sweetcorn, wheat, cereals, pasture or forage brassicas within 42 days of plating and grass seed oversown into pasture (stitching) within 21 days from sowing

® Registered Trademark of BASF

© Copyright 2019, BASF New Zealand Limited