



Stomp[®] Xtra

For the control of annual grasses and broadleaf weeds in certain fruit crops and onions.

Contains 455 g/litre pendimethalin in the form of a capsulated suspension.

WARNING

Keep out of reach of children

NET CONTENTS: 10 Litres

Registered to and distributed by:
BASF New Zealand Limited
P.O. Box 407, Auckland 1140
Phone: 0800 932 273
Fax: 0800 940 000



© Registered Trademark of BASF

READ LABEL COMPLETELY BEFORE OPENING OR USING

Hazard Classification

6.9B, 9.1A, 9.2A



WARNING

HARMFUL. May cause organ damage from repeated oral exposure at high doses.
VERY TOXIC TO AQUATIC LIFE WITH LONG LASTING EFFECTS. May cause long-term adverse effects in the aquatic environment. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites.
VERY TOXIC TO THE SOIL ENVIRONMENT. Selective herbicide - very toxic to some plant species (certain plants may be killed or damaged from uptake of this product).

PRECAUTIONS

HANDLING: This product must be under the personal control of an APPROVED HANDLER when used in a wide dispersive manner or by a commercial contractor.
STORAGE: Store in original container, tightly closed, in a dry, cool, well-ventilated place, out of direct sunlight and away from food and animal feeds. Stores containing 100 litres or more of this product require emergency response plans and secondary containment systems and are subject to signage.
TRANSPORT: Do NOT carry this product on a passenger service vehicle.
SPILLAGE: Prevent further spillage and product from entering waterways. Wear overalls, suitable gloves, facemask or goggles. Prevent further spillage. Take up spilled product with An inert absorbant material (e.g. vermiculite) and place in sealable container for disposal. Wash down affected area with water plus detergent. Absorb and collect washings and place in the same sealable container. Dispose of the waste safely in suitable landfill.
PERSONAL PROTECTION: When mixing and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, suitable gloves, and face shield or safety goggles.
GENERAL SAFETY AND HYGIENE MEASURES: Avoid contact with skin, eyes and clothing. Avoid inhalation of vapour, mist or spray. Do NOT eat, drink or smoke while using this product. Remove protective clothing and wash hands and exposed skin thoroughly with soap and water before meals and after work. Wash outside of gloves before removing. Wash protective clothing immediately after use. Store and wash work clothing separately.
USE: Apply with well-maintained and accurately calibrated equipment. Do NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do NOT spray over, or allow drift over, surface water. Do NOT apply STOMP XTRA under adverse weather conditions, or with spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, croplands or pastures. Clean out application equipment thoroughly after use. Do NOT contaminate water when disposing of equipment wash waters.
RECORD KEEPING: Records of use must be kept if 3 litres or more is applied in a place where the substance is likely to enter air or water and leave the place.
NOTE: Storage, application and record keeping must be as described in the current version of the New Zealand Standard for the Management of Agrichemicals NZS8409.
DISPOSAL: Container: Triple rinse empty container and add rinsate to the spray tank. Recycle the rinsed container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Product: Dispose of this product only by using according to the label or at an approved landfill. Do NOT burn product. Do NOT contaminate surface water with chemical or empty container.

FIRST AID

IF SWALLOWED: Do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISONS (0800 764 766) or a doctor if you feel unwell. Rinse mouth. **IF ON SKIN OR HAIR:** Remove contaminated clothing and flush skin and hair with running water. Do NOT scrub the skin. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. **IF INHALED:** Remove to fresh air and keep at rest in a position comfortable for breathing. Get medical advice/attention if you feel unwell.

GENERAL INFORMATION

STOMP XTRA is a selective herbicide recommended for use in certain fruit crops and onions. DO NOT USE ON SWEETCORN OR CROPS GROWN FOR SEED. Following application of STOMP XTRA to the soil and its absorption by the plant, both root and shoot growth are inhibited in susceptible plants. Affected plants die shortly after germination or following emergence from the soil. STOMP XTRA will not control perennial or established weeds. Weed control with STOMP XTRA may be reduced where soil organic matter exceeds 6%. Approximately 12 mm of rainfall or irrigation within ten days of application is required for optimum performance. Reduced weed control can be expected when used under dry soil conditions.

FOLLOWING CROP RESTRICTIONS

Under prolonged dry conditions or prolonged cold wet conditions certain crops are susceptible to injury when planted in soil previously treated with STOMP XTRA. Do not plant sensitive grasses (such as ryegrass), cucurbits or beets for at least 6 months following STOMP XTRA application.

WEEDS CONTROLLED BY STOMP XTRA

Black nightshade, Chickweed, Cornbind*, Fathen, Field pansy, Fumitory, Henbit, Redroot*, Scarlet pimpernel, Shepherds purse*, Speedwell, Spurrey, Stinking mayweed*, Storksbill*, Wild oats*, Wireweed.

* These weeds controlled under favourable conditions or by using higher rates

DIRECTIONS FOR USE

Crop	Rate	Critical Comments
Fruit Crops Citrus, Feijoas, Grapes (dormant only), Kiwifruit, Pipfruit (including Nashi), Stonefruit, Tamarillos	3.6-4.4 litres/ha	Apply STOMP XTRA as a bare ground treatment. On established plantings where simazine safety has been proven simazine may be added as a tank mix at recommended rates. STOMP XTRA may be used on one year old trees in the year of planting. Avoid spray contact with green bark or drift on to green buds or foliage. STOMP XTRA may be mixed with knockdown herbicides if small weeds are present but should otherwise be applied to bare earth. Irrigation or rain should follow within 10 days of application or weed control may be reduced.
Onions	0.75 - 1.1 litres/ha 1.1-1.5 litres/ha	PRE CROP EMERGENCE Apply STOMP XTRA as soon as possible after sowing. Do NOT apply later than 4 days after sowing. Application close to crop emergence will result in crop damage. Use the higher rate on soils high in organic matter or clay content. Do NOT use on coarse open soils, particularly sandy soils. POST CROP EMERGENCE Apply STOMP XTRA from the 2 true leaf stage. A further application may be made 2-4 weeks later but not after the 4 true leaf stage. Post-emergence applications with STOMP XTRA will not control established weeds but will extend the period of residual weed control.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

SOIL PREPARATION (Seeded and transplanted crops)

Prepare the seedbed with a fine, firm tilth free of ridges, clods or trash. The seed should be sown so that after consolidation it is covered by a minimum of 15 mm of settled soil for onions. Do NOT disturb the soil by rolling or harrowing after STOMP XTRA has been applied.

MIXING

Fill the spray tank to one quarter capacity with water with the agitation system operating. If required, add a tank mix partner at this stage and mix thoroughly. Fill the spray tank to one half capacity and add the required amount of STOMP XTRA and the rest of the water, mixing thoroughly. Agitation is required during spraying and after a stoppage.

APPLICATION

Use a conventional sprayer with either mechanical or bypass agitation. Apply the recommended amount of STOMP XTRA herbicide in sufficient water to ensure accurate cover. This should be from 150-400 litres of water per hectare.

CONDITIONS OF SUPPLY

Read conditions of supply before buying and using. If these conditions are unacceptable, return at once unopened. All goods supplied by BASF New Zealand Limited are of a high grade and conform to the chemical description of the label. However, to the extent permitted by law, neither BASF nor any of its Distributors accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential, howsoever arising, in connection with the supply of those goods including without limitation their storage, application, handling or use. BASF New Zealand Limited, nor any of its Distributors, make no warranties of any kind, express or implied, with respect to fitness for purpose (including crop injury, ineffectiveness or any other unintended consequence), merchantable quality or any other matter with respect to these goods. When you are a consumer as defined in the Consumer Guarantees Act 1993 (the "Act") these conditions will be subject to the provisions of the Act. However the Act will not apply to goods supplied to you for business purposes.

Approved pursuant to the HSNO Act 1996, Code HSR000547.

See www.epa.govt.nz for approval controls.

Registered pursuant to the ACVM Act 1997, No. P5921.

See www.foodsafety.govt.nz for registration conditions.

Date of Manufacture:

Batch No.:

EMERGENCY PHONELINK
24 Hour Emergency Advice in an Emergency Only
0800 944 955

© Copyright 2013, BASF New Zealand Limited