

# Belanty®

## Fungicide

### Belanty Quick Fire Fact Sheet APPLES AND PEARS

**BASF**  
We create chemistry

<b>Q</b>	<b>Pipfruit diseases controlled?</b>	<b>A</b>	Black spot in apples and pears and powdery mildew in apples
<b>Q</b>	<b>Product type?</b>	<b>A</b>	Systemic fungicide with protectant and curative activity
<b>Q</b>	<b>Mode-of-Action group?</b>	<b>A</b>	FRAC Group 3, Demethylation Inhibitors (DMI)
<b>Q</b>	<b>Active ingredient?</b>	<b>A</b>	Mefentrifluconazole
<b>Q</b>	<b>Formulation?</b>	<b>A</b>	75 g/L suspension concentrate
<b>Q</b>	<b>Pack size?</b>	<b>A</b>	10 L plastic Ecopack (eligible for Agrecovery)
<b>Q</b>	<b>Application rate?</b>	<b>A</b>	80 mL/100 L, full dilute application. For concentrate spraying adjust rate accordingly. Do not apply more than 1.6 L/ha per application.
<b>Q</b>	<b>Adjuvant?</b>	<b>A</b>	Not required
<b>Q</b>	<b>Rainfastness?</b>	<b>A</b>	Once dry on the crop
<b>Q</b>	<b>Compatibility?</b>	<b>A</b>	Compatible with most commonly used fungicides, insecticides and plant growth regulators
<b>Q</b>	<b>Tank mixing?</b>	<b>A</b>	Always tank mix with a fungicide from another mode of action group
<b>Q</b>	<b>Resistance management?</b>	<b>A</b>	Do not make more than two consecutive applications or apply more than four times per season
<b>Q</b>	<b>Withholding period?</b>	<b>A</b>	Apply no later than 80% petal fall
<b>Q</b>	<b>Grazing?</b>	<b>A</b>	Do not allow stock to graze within treated orchards until after harvest (no slaughter withholding period applies)
<b>Q</b>	<b>Buffer zones?</b>	<b>A</b>	5 m from waterbodies (30 m downwind)

For more information on Belanty, visit [crop-solutions.basf.co.nz](https://crop-solutions.basf.co.nz) or call **0800 932 273**

**ALWAYS READ AND FOLLOW LABEL DIRECTIONS BEFORE USING ANY PRODUCT IN THIS FACT SHEET.**

© Copyright BASF 2021. Sercadis® and Pristine® are registered trademarks of BASF.

# Belanty®

## Fungicide

**BASF**  
We create chemistry

### Belanty Quick Fire Fact Sheet GRAPES

<b>Q</b>	Grape diseases controlled?	<b>A</b>	Powdery mildew
<b>Q</b>	Product type?	<b>A</b>	Systemic fungicide with protectant and curative activity
<b>Q</b>	Mode-of-Action group?	<b>A</b>	FRAC Group 3, Demethylation Inhibitors
<b>Q</b>	Active ingredient?	<b>A</b>	Mefentrifluconazole
<b>Q</b>	Formulation?	<b>A</b>	75 g/L suspension concentrate
<b>Q</b>	Pack size?	<b>A</b>	10 L plastic Ecopack (eligible for Agrecovery)
<b>Q</b>	Application rate?	<b>A</b>	80 mL/100 L, full dilute application. For concentrate spraying adjust rate accordingly. Do not apply more than 800 mL/ha per application.
<b>Q</b>	Adjuvant?	<b>A</b>	Not required
<b>Q</b>	Rainfastness?	<b>A</b>	Once dry on the crop
<b>Q</b>	Compatibility?	<b>A</b>	Compatible with most commonly used fungicides, insecticides and plant growth regulators
<b>Q</b>	Tank mixing?	<b>A</b>	Always tank mix with a fungicide from another mode of action group or alternate groups between applications
<b>Q</b>	Resistance management ?	<b>A</b>	Do not use more than two Belanty applications per season
<b>Q</b>	Withholding period?	<b>A</b>	Apply no later than pre-bunch closure (E-L 31)
<b>Q</b>	Leaf plucking?	<b>A</b>	A 6-month slaughter withholding period must be observed after stock have been removed from the vineyard.
<b>Q</b>	Grazing?	<b>A</b>	Do not allow stock to graze within treated vineyards until after harvest

For more information on Belanty, visit [crop-solutions.basf.co.nz](http://crop-solutions.basf.co.nz) or call **0800 932 273**

**ALWAYS READ AND FOLLOW LABEL DIRECTIONS BEFORE USING ANY PRODUCT IN THIS FACT SHEET.**

© Copyright BASF 2021.