

WARNING
KEEP OUT OF REACH OF CHILDREN
READ LABEL BEFORE USE

IMTREX[®]

Contains 62.5 g/litre fluxapyroxad in the form of an emulsifiable concentrate

GROUP 7 FUNGICIDE

Systemic cereal fungicide for the control of diseases in barley and wheat crops

Contents: 10 L

Registered to and distributed by:

BASF New Zealand Limited

NZBN 9429032151299

P.O. Box 407, Auckland 1140

Phone: 0800 932 273

crop-solutions.basf.co.nz

Hazard Classification

6.1E, 6.1D, 6.4A, 6.9B, 9.1D, 9.2C, 9.3C



WARNING

May be harmful if swallowed. Harmful if inhaled. Causes serious eye irritation. May cause damage to organs through prolonged or repeated exposure. Toxic to aquatic life with long lasting effects. Harmful to the soil environment. Harmful to terrestrial vertebrates.

PRECAUTIONS

HANDLING: Obtain special instructions before use. DO NOT handle until all safety precautions have been read and understood. Use personal protective equipment as required.

STORAGE: Keep out of reach of children. Store locked up. Store in original container, tightly closed, in a locked, dry, cool, well-ventilated place away from food and animal feeds. Keep it dry and protect from frost. Keep away from heat and protect from direct sunlight.

TRANSPORT: DO NOT carry this product on a passenger service vehicle.

SPILLAGE: Prevent further spillage and product from entering waterways. Collect spillage. Absorb spill with an inert material (e.g. vermiculite) and place in a labelled, sealable container for disposal. Wash affected area with water and detergent. Absorb and collect washings and place in the same sealable container as the spilled material. Dispose of the waste safely in an approved landfill.

PERSONAL PROTECTION: Wear suitable protective clothing, suitable protective gloves and eye/face protection when applying spray. Wear suitable protective clothing (coveralls), suitable protective gloves and face protection (face shield) when preparing spray.

GENERAL SAFETY AND HYGIENE MEASURES: Avoid contact with skin, eyes and clothing. DO NOT breath vapour or spray mist. DO NOT eat, drink or smoke when using this product. Wash hands and exposed skin thoroughly after handling. Contaminated clothing should not be allowed out of the workplace. Wash protective clothing immediately after use. Store work clothing separately. Use only outdoors or in well-ventilated area.

USE AND ENVIRONMENTAL PROTECTION: Apply with well-maintained and accurately calibrated equipment. Clean out application equipment thoroughly after use. DO NOT apply directly into or onto water. Take all reasonable steps to ensure that the product does not cause any significant adverse effects to the environment beyond the application area. Apply this substance at a maximum application rate of 1.25 L/ha in a maximum two applications per year and with a minimum application interval of 21 days. DO NOT allow drift over surface water such as ponds, waterways or drains. DO NOT clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

DISPOSAL: Container: DO NOT reuse the container for any other purpose. Triple rinse empty container and add rinsate to the spray tank. Recycle through Agrecovery (0800 247 326, www.agrecovery.co.nz). Product: Dispose of this product only by using according to the label or at an approved landfill or at an approved facility. DO NOT burn product.

NOTE: Storage and application must be as described in the current version of the New Zealand Standard for the Management of Agrichemicals NZS8409.

FIRST AID

If exposed or concerned get medical advice/attention. If medical advice is needed, have the product label at hand.

IF SWALLOWED: DO NOT induce vomiting. Rinse mouth. Call the National Poisons Centre 0800 POISONS (0800 764 766) or a doctor if you feel unwell. Have product label on hand. **IF ON SKIN:** Remove contaminated clothing and wash skin with plenty of soap and water. DO NOT scrub the skin. If skin irritation or rash occurs, get medical advice/attention. Wash contaminated clothing before reuse. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice/attention. **IF INHALED:** Remove to fresh air and keep at rest in a position comfortable for breathing. Call a POISON CENTRE or a doctor/physician. Get medical advice/attention if you feel unwell.

GENERAL INFORMATION

IMTREX is a broad spectrum contact and systemic fungicide with protectant and eradicant activity. IMTREX is taken up via the stem and foliage and is translocated upwards and outwards to provide some protection of new growth. Thorough coverage of the crop is necessary for best results. IMTREX has curative activity against early stages of infection but optimum results are obtained from application prior to the onset of infection.

RESISTANCE MANAGEMENT

IMTREX contains fluxapyroxad, a carboxamide fungicide from the SDHI (Succinate Dehydrogenase inhibitors) mode of action group **GROUP 7 FUNGICIDE**. Resistance to this fungicide could develop from excessive use. To minimize this risk use strictly in accordance with label instructions and resistance management strategies that exist for any fungus listed on the label. Frequency of application must be strictly adhered to in accordance with the Directions for Use. IMTREX should be used in a programme that incorporates fungicides with different modes of action to that of GROUP 7. Apply no more than two IMTREX sprays per season. Refer to the NZCPR web site < www.resistance.nzpps.org > for more detailed information.

DIRECTIONS FOR USE

It is an offence to use this product on animals.

CROP	DISEASE	REMARKS
Rate of Use: 1.25 litres/ha IMTREX in a minimum of 150 litres of water/ha by ground or 60 litres/ha by air. IMTREX must be tank mixed with 625 mL/ha Opus or an approved triazole-based product that is also recommended for the same target diseases in barley and wheat. The triazole-based tank mix product should be applied at the recommended label rate.		
Barley	Leaf rust Net blotch (spot & net forms) Scald Powdery mildew Ramularia leaf and awn spot	Apply as a preventative spray on susceptible cultivars and when conditions favour disease. Otherwise apply at the first appearance of disease. Continued disease pressure or reinfection may require a further treatment 3-5 weeks later. Apply up to the end of flowering (Zadok GS 69).
Wheat	Brown rust Powdery mildew Speckled leaf blotch Stripe rust	Apply as a preventative spray on susceptible cultivars and where conditions favour disease. Otherwise apply at the first appearance of disease. Continued disease pressure or reinfection may require a further treatment 3-5 weeks later. Apply up to the end of flowering (Zadok GS 69).
	Glume blotch	Apply as a preventative spray around flag leaf emergence. For continued protection of the ear a further treatment may be applied up to the end of flowering (Zadok GS 69).

Mixing:

Add the required amount of IMTREX and then the approved triazole-based formulation directly into the partly filled spray tank with agitation operating. Complete filling and maintain agitation until spraying is complete. DO NOT leave spray mix standing overnight in the spray tank.

Compatibility:

IMTREX may be tank mixed with most commonly used fungicides (e.g. Comet[®], OPUS[®] and other triazole-based formulations, including epoxiconazole- or prothioconazole-based formulations), insecticides, herbicides (e.g. fluroxypyr) and growth regulators (e.g. Cycocel[®] or Terpal[®]). All tank mixes should be used **immediately** after preparation.

Rainfall:

DO NOT apply if rainfall is likely before IMTREX dries on the crop.

Withholding Periods:

Cereal green feed/silage: Allow 28 days between last application and introduction of stock for grazing or cutting for silage.

Cereal grain: DO NOT harvest within 42 days of last application.

Note: observe the withholding period of tank mix partners

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

CONDITIONS OF SUPPLY

Read conditions of supply before buying and using. If these conditions are unacceptable, return at once unopened. All goods supplied by BASF New Zealand Limited are of a high grade and conform to the chemical description of the label. However, to the extent permitted by law, neither BASF nor any of its Distributors accept any liability or responsibility whatsoever for any loss, damage or injury whether direct, indirect or consequential howsoever arising in connection with the supply of those goods including without limitation their storage, application, handling or use. BASF New Zealand Limited nor any of its Distributors, make no warranties of any kind, express or implied, with respect to fitness for purpose (including crop injury, ineffectiveness or any other unintended consequence), merchantable quality or any other matter with respect to these goods. When you are a consumer as defined in the Consumer Guarantees Act 1993 (the "Act") these conditions will be subject to the provisions of the Act. However, the Act will not apply to goods supplied to you for business purposes.

Registered pursuant to the ACVM Act 1997, No. P9613

See www.foodsafety.govt.nz for registration conditions

Approved pursuant to the HSNO Act 1996, Code HSR101303

See www.epa.govt.nz for approval controls

Batch No:

Date of Manufacture:

Registered to and distributed by:

BASF New Zealand Limited

NZBN 9429032151299

P.O. Box 407, Auckland 1140

Phone: 0800 932 273

crop-solutions.basf.co.nz

EMERGENCY PHONELINK
24 Hour Advice in an Emergency Only
0800 944 955



IIIIII BAR CODE IIIIII

® Registered Trademark of BASF

© Copyright 2019, BASF New Zealand Limited