



Q	Pipfruit diseases controlled?	A	Black spot in apples and pears and powdery mildew in apples
Q	Product type?	A	Systemic fungicide with protectant and curative activity
Q	Mode-of-Action group?	A	FRAC Group 3, Demethylation Inhibitors (DMI)
Q	Active ingredient?	A	Mefentrifluconazole
Q	Formulation?	A	75 g/L suspension concentrate
Q	Pack size?	A	10 L plastic Ecopack (eligible for Agrecovery)
Q	Application rate?	A	80 mL/100 L, full dilute application. For concentrate spraying adjust rate accordingly.
Q	Adjuvant?	A	Not required
Q	Rainfastness?	A	Once dry on the crop
Q	Compatibility?	A	Compatible with most commonly used fungicides, insecticides and plant growth regulators
Q	Tank mixing?	A	Always tank mix with a fungicide from another mode of action group
Q	Resistance management?	A	Do not make more than two consecutive applications or apply more than four times per season
Q	Withholding period?	A	Apply up to 35 days pre-harvest (check with your exporter). For nil detectable residues apply no later than 80% petal fall.
Q	Grazing?	A	Do not allow stock to graze within treated orchards until after harvest (no slaughter withholding period applies)
Q	Buffer zones?	A	5 m from waterbodies (30 m downwind)

For more information on Belanty, visit crop-solutions.basf.co.nz or call 0800 932 273

## Belanty **D-BASF** Fungicide We create chemistry **Belanty Quick Fire Fact Sheet** GRAPES

Q	Grape diseases controlled?	A	Powdery mildew
Q	Product type?	A	Systemic fungicide with protectant and curative activity
Q	Mode-of-Action group?	A	FRAC Group 3, Demethylation Inhibitors
Q	Active ingredient?	A	Mefentrifluconazole
Q	Formulation?	A	75 g/L suspension concentrate
Q	Pack size?	A	10 L plastic Ecopack (eligible for Agrecovery)
Q	Application rate?	A	80 mL/100 L, full dilute application. For concentrate spraying adjust rate accordingly.
Q	Adjuvant?	A	Not required
Q	Rainfastness?	A	Once dry on the crop
Q	Compatibility?	A	Compatible with most commonly used fungicides, insecticides and plant growth regulators
Q	Tank mixing?	A	Always tank mix with a fungicide from another mode of action group or alternate groups between applications
Q	Resistance management?	A	Do not use more than two Belanty applications per season (check with your exporter)
Q	Withholding period?	A	Apply no later than pre-bunch closure (E-L 31) (nil detectable residues in wine)
Q	Leaf plucking?	A	A 6-month slaughter withholding period must be observed after stock have been removed from the vineyard
Q	Grazing?	A	Do not allow stock to graze within treated vineyards until after harvest

For more information on Belanty, visit crop-solutions.basf.co.nz or call 0800 932 273

