

Revystar[®]

Fungicide

FACT SHEET

A new first choice for foliar disease in cereals

Revystar[®] is an important dual-active foliar fungicide that controls the key diseases of cereals. Adding Revystar to your spray rotations will give you superior control of *Ramularia* leaf spot in barley and help maintain excellent control of diseases like speckled leaf blotch in wheat by reducing resistance pressure.

Crop registrations:

Wheat, barley, triticale, ryecorn and oats

Active ingredients:

Mefentrifluconazole 100 g/L
Fluxapyroxad 50 g/L

Pack size: 10 L - Agrecovery*

Diseases controlled:

Barley: *Ramularia* leaf spot
Scald
Brown rust

Oats: Crown rust

Ryecorn: Leaf (brown) rust and scald

Triticale: Leaf (brown) and stripe rust

Wheat: Speckled leaf blotch (*Septoria tritici* blotch)
Stripe rust
Leaf rust
Powdery mildew

Rates:

1.5 L/ha Revystar in at least:

- 150 L/ha water for ground application
- 60 L/ha water for aerial application

WHPs:

Grain & straw stubble: DO NOT harvest or graze within 35 days of last application

Green feed/silage: allow 28 days between last application and introduction of stock for grazing or cutting for silage

- Outstanding control of *Ramularia* in barley and excellent protection against other key diseases creating yield potential in cereal crops.
- Extra green leaf retention, which also promotes higher yields.
- The unique combination of the most powerful Group 3 triazole fungicide and market-leading Group 7 SDHI chemistry provides excellent control of pathogens with resistance to other fungicide modes of action.
- Less reliance on favourable weather for application and performance thanks to superior rainfastness and tolerance of cool temperatures.
- The added convenience of having a single very effective solution for barley, wheat and other cereal crops.
- Introducing Revystar's unique mode of action to control speckled leaf blotch in wheat will help maintain and extend the effectiveness of older chemistry.



*Containers are eligible for Agrecovery recycling.

BASF
We create chemistry


How and when to use Revystar

T2 is the 'king hit' application timing for all cereal crops

AUTUMN BARLEY

Optimal *Ramularia* timings:

T1 (GS 31–33) and T2 (GS 55–59), treating the top 4 leaves & ear. 2 sprays will usually be needed, at T2 (GS 55–59) and 21–28 days earlier. Adding a multisite is beneficial at T1.




SPRING BARLEY

Optimal *Ramularia* timing:

T2 (GS 49). An earlier Revystar spray may be an option when disease pressure is high.

In whole crop silage, a single Revystar application in T1-T2 will maintain green leaf area and boost grain fill.




WHEAT

Optimal SLB timings:

T1 (GS 31–33) and T2 (GS 39–45), treating the top 4 leaves. Apply Revystar twice if disease pressure is high. If not, save Revystar for the critical T2 application.

In all crops, use an alternative triazole +/- multisite at other timings.



Resistance management

Revystar should be used in a programme that incorporates fungicides with different modes of action to that of Group 3 and Group 7 fungicides. Apply no more than two Revystar sprays per season. Refer to the NZCPR web site www.resistance.nzpps.org for more detailed information.

For more information about Revystar, visit crop-solutions.basf.co.nz

or call **0800 932 273** or your local territory manager:

Canterbury & West Coast **Colin Dunstan 027 222 7205**

Otago & Southland **Geoff Booth 021 744 025**

Upper North Island **Brendan Arnet 027 336 5544**

Lower North Island **Tim Geuze 021 730 438**

BASF
We create chemistry

ALWAYS READ AND FOLLOW LABEL DIRECTIONS BEFORE USING ANY PRODUCT

© Copyright BASF 2024. ® Registered trademarks of BASF. *Registered trademark. ACVM Registration numbers: Revystar: P009654, Comet: P006017, Opus: P004365

BASFNZ0057 09.24