

Superior disease management starts with Revylution[®]. It delivers exceptional control of *Septoria tritici* blotch (STB) and leaf rust - as well as doing a great job on stripe rust in wheat. It has quickly established itself as the '**must-have**' fungicide at the critical T2 timing in wheat.



Crop registrations:

Wheat Barley

Diseases controlled:

Wheat: Leaf rust

Septoria tritici blotch

Stripe rust

Barley: Brown rust

Ramularia leaf spot

Rates:

1.5 L/ha Revylution in at least:

- 150 L/ha water for ground application
- 60 L/ha water for aerial application

WHPs:

Cereal grain, straw and stubble:

DO NOT harvest or graze within 42 days of last application

Cereal green feed/silage: allow 28 days between last application and introduction of stock for grazing or cutting for silage

- The 'must-have' DMI in your tank mix
- A highly flexible mix partner for targeted disease control
- Powered by a unique DMI that works where others are beginning to fail
- Ultra-rapid absorption delivers exceptionally reliable performance and superior rainfastness
- Extended protection for weeks after application
- Highly flexible and reliable tank-mix partner
- 100 x more powerful against *Septoria tritici* blotch even when resistance has compromised older DMIs






Revylution®

Fungicide

How and when to use Revylution

Revylution can be applied as a preventative spray on susceptible cultivars when conditions favour disease or at the first appearance of disease. Continued disease pressure may require a further treatment 3 to 5 weeks later. Do not apply after the end of flowering (Zadok GS 69).

T2 is the 'king hit' application timing for SLB in wheat



The optimal timings for *Septoria tritic*i blotch control are T1 (GS 31–33) and T2 (GS 39–45), treating the top 4 leaves. Apply Revylution twice if disease pressure is high. If not, save Revylution for the critical T2 application.

Compatibility

Revylution is compatible with most commonly used fungicides (e.g. Comet[®], SDHIs, folpet) insecticides, herbicides (e.g. Granstar*, Starane* Xtra, or Duplosan* KV) and growth regulators (e.g. Cycocel[®], Moddus* or Terpal[®]).

Resistance management

Revylution should be used in a programme that incorporates fungicides with different modes of action to that of DMI (Group 3 fungicides). Apply no more than three DMI fungicide sprays per season. Refer to the NZCPR website **www.resistance.nzpps.org** for more detailed information.

For more information about Revylution®, visit **crop-solutions.basf.co.nz** or call **0800 558 399** or your local territory manager:

Northland

Molly Green | 027 336 5544

Canterbury & West Coast

Colin Dunstan | 027 222 7205

Otago & Southland

Simon Von Tunzelman | 027 237 4402

