



Take control of *Septoria tritici* blotch along with other key diseases in wheat

With its superior, targeted control of *Septoria tritici* blotch, including strains that are less susceptible to demethylation inhibitors (DMI) fungicides, Revylution® will help simplify your fungicide rotations and deliver extra value for money.

Q Why should I start using Revylution?

A This fungicide is the first isopropanol-azole active ingredient, with a uniquely flexible molecule that can adapt to mutations in the target pathogens and control strains of disease that are less susceptible to other DMI fungicides.

Revylution is your flexible 'must-have' tank-mix, delivering exceptional disease control where other DMIs are beginning to fail.

Q Which diseases does Revylution control?

A Wheat

- *Septoria tritici* blotch
- Leaf rust
- Stripe rust

Barley

- Ramularia leaf spot
- Brown rust

Q What's the best application timing for Revylution against *Septoria tritici* blotch?

A For *Septoria tritici* blotch control the king hit is T2: when the flag leaf is fully emerged (GS 39) to booting (GS 45). In severe conditions an earlier application at GS 31-32 may be required.

The best timing for leaf rust control is around the booting and end of flowering stage (GS45-69).

Q What if I need to control Ramularia in my barley?

A Revylution delivers outstanding control of Ramularia in barley and T2 is your optimal timing window to maximise your returns.



BASF
We create chemistry



Revylution®

Fungicide

Q What is Revylution compatible with?

A Revylution is the ideal mixing partner for a range of complementary fungicide options including Qol's such as Comet®; SDHIs and other MoAs such as folpet.

Revylution also provides you with the flexibility to mix with insecticides, and herbicides such as Granstar*, Starane* Xtra or Duplosan* KV; and growth regulators such as Cycocel®, Moddus* or Terpal®.

Check the manufacturers' recommendations for rates of any tank-mix product.

Q How will Revylution strengthen my resistance management program?

A Revylution sets a new standard for resistance management, delivering unmatched control particularly where older DMIs are beginning to fail.

Revylution, brings a more robust Group 3 triazole to the market with its flexible, iso-propanol neck, ensuring on-going efficacy as populations continue to shift in sensitivity to older triazoles.

Do not apply Group 3 triazole foliar fungicides more than three times per season

Q What are the withholding periods?

A Cereal grain, straw and stubble: DO NOT harvest within 42 days of last application.

Cereal green feed/silage: Allow 28 days between last application and introduction of stock for grazing or cutting for silage.

For tank-mixing, observe the longest WHP of any tank-mix partner.

Revylution product profile

Active ingredient	Revysol® (mefentrifluconazole) 100 g/L
Mode of action	Group 3 – Demethylation inhibitors (DMI)
Pack size	10 L
Adjuvant	Not required
Rainfastness	Once dry on the crop
Application rate	1.5 L/ha
Water rates	Minimum of 150 L/ha by ground or 60 L/ha by air
Directions of use	Add required amount of Revylution directly into partly filled spray tank with agitation operating. Complete filling and maintain agitation until spraying complete. DO NOT leave spray mix standing overnight.

For more information about Revylution®, visit **crop-solutions.basf.co.nz** or call **0800 558 399** or your local territory manager:

Northland

Molly Green | 027 336 5544

Canterbury & West Coast

Colin Dunstan | 027 222 7205

Otago & Southland

Simon Von Tunzelman | 027 237 4402

BASF
We create chemistry

ALWAYS READ AND FOLLOW LABEL DIRECTIONS BEFORE USING ANY PRODUCT IN THIS Q&A SHEET.

© Copyright BASF 2025. ® Registered trademarks of BASF. *Registered trademark. ACVM registration numbers: Revylution: P009923, Comet: P00617, Cycocel: P001493 and Terpal: 3483
AU55-P00001950 0825