



For superior control of broadleaf & grass weeds

Eliminate yield sapping weeds from your maize and sweetcorn crops with the powerful in-crop protection of Arietta® herbicide. This post emergent herbicide can be tank-mixed to further extend the weed spectrum and delivers the proven reliability growers have come to expect.

Q Why should I use Arietta?

A Arietta controls broadleaf weeds (up to 8 leaves) and grass weeds (up to 3 tillers). It's proven performance and tank-mix flexibility has elevated Arietta to the top of the table in post-emergent weed control.

Q What are the recommended application rates?

A Arietta 200 mL/ha + atrazine 500 g ai/ha + ECO** 500 mL/100 L of water

Q What are some of the tank-mixing options for maize to extend weed control?

A Atrazine - Enhances Arietta performance on labelled weeds

Nicosulfuron (WG formulations only) - Ryegrass and couch (twitch)

Clopyralid - Californian thistle, yarrow and volunteer potatoes

Dicamba - Californian thistle, field bindweed (*Convolvulus arvensis*) and cornbind (*Polygonum convolvulus*)

PLEASE NOTE: Do not use dicamba, clopyralid or nicosulfuron mixtures on sweetcorn

Q What oil/adjuvant should I use?

A An ECO is recommended as they provide Arietta with the most effective penetrant system, e.g. Hasten™, Kwickin™ or Bonza® Gold

Q What is the recommended water rate?

A Minimum of 200 L/ha

"Coverage is Key" – for the best results use 300-400 L/ha



Q Rainfastness?

A Do not apply if rain is expected within 2 hours of application

Post-emergent broadleaf and grass weed control in maize and sweetcorn



Broadleaf weeds (up to 8 leaves)			Grass weeds (up to 3 tillers)
Apple of Peru	Fathen	Velvetleaf (before it lignifies)	Barnyard grass
Black nightshade	Fleabane	White clover	Smooth witchgrass
Broad-leaved dock (seedling)	Groundsel	Willow weed	Summer grass
Chamomile	Large flowered mallow		Yellow bristle grass
Chickweed	Scrambling speedwell		
Dandelion	Twin cress		

The BASF portfolio fit in the maize and sweetcorn programme

Soil & planting preparation	Crop establishment (pre-emergence)	Vegetative before canopy closure	Reproductive	Harvest & post harvest
 Sharpen® 25 g/ha Herbicide + glyphosate Herbicide ECO + 500 mL/100 L water Adjuvant	 Sharpen 150 g/ha Herbicide + acetochlor 3 L/ha Herbicide	 Arietta 200 mL/ha Herbicide + atrazine 500 g ai/ha Herbicide ECO + 500 mL/100 L water Adjuvant	 Opus® 1 L/ha Fungicide + Comet® 1 L/ha Fungicide Timing V5-VT Control of rust	

Broadleaf control Arietta + atrazine + ECO

Grass weed control Arietta + atrazine + ECO

Mallow	Black nightshade	Barnyard grass (tillering)	Barnyard grass (seedling)	
				At application
				4 days after treatment
				9 days after treatment



Scan the QR code to find out more about **Arietta Herbicide** or call **0800 932 273**

ALWAYS READ AND FOLLOW LABEL DIRECTIONS.

This leaflet is intended as general advice. Arietta ACVM Reg. No. P008825, Sharpen ACVM Reg. No. P008391, Opus ACVM Reg. No. P004365 and Comet ACVM Reg. No. P006017 are registered trademarks of BASF. Hasten™ is a Trademark of the Victorian Chemical Company. Bonza Gold® is a registered trademark of Nufarm, registered pursuant to the ACVM Act 1997, No. P8871 © Copyright BASF 2025 ® Registered trademark of BASF * Registered trademark. AU55-P00002116 0825

BASF
We create chemistry