

Sharpen®

Herbicide

Powerful pre-emergence control of key broadleaf weeds in maize & sweet corn crops

Sharpen® is the only Group 14 herbicide registered for use in maize, delivering excellent control whilst also helping growers manage their herbicide resistance management strategies. Using Sharpen in a tank-mix with acetochlor for pre-emergent applications straight after sowing provides very effective control of broadleaf weeds, including weeds that are resistant to triazine herbicides (e.g. atrazine).



- Excellent control of major broadleaf weeds, including triazine-resistant willow weed, nightshade and fathen.
- Tank-mixes with pre-emergence grassweed herbicides broaden their spectrum of activity.
- Sharpen is from a different herbicide mode of action group strengthening herbicide resistance management in maize and sweet corn crops.
- Extended residual activity in the soil.
- No need to decontaminate the spray rig, unlike other PPO herbicides.
- Favourable environmental, toxicological and ecotoxicological profiles.

Mode of action

Group 14 Protoporphyrinogen-oxidase inhibitors (PPOs)

Pre-emergence crop registrations

Maize
Sweet corn

WHP

Do not graze or cut for feed within 28 days after application

Up to 8 weeks' pre-emergent control of:

Fathen
Black nightshade
Broadleaf docks (seedling)
Willow weed
Wireweed

Up to 8 weeks' pre-emergent suppression of:

Pink shamrock (oxalis)

Sharpen®

Herbicide

How and when to use Sharpen for pre-emergence control

Rate

Sharpen 150 g/ha + acetochlor, S-metolachlor or Frontier-P® at label rates.

No adjuvant is required for this use pattern.

Timing

Apply as soon as possible after planting and before the crop or weeds emerge.

Optimum broadleaf weed control and resistance management

Substituting Sharpen for atrazine as a pre-emergence herbicide needn't mean not using atrazine at all. This suggested program incorporates atrazine in a tank-mix with Arietta® for post-emergence control to follow up the extended residual pre-emergence activity of Sharpen mixed with acetochlor.

Both Sharpen and Arietta (mode of action Group 27) are in the moderate risk category for developing resistance and provide excellent control of key weeds. Arietta is the only herbicide registered to control velvetleaf in maize.

This suggested program uses herbicides from 5 different herbicide groups, thereby maximising resistance management.

Suggested tank-mixes and application timings



* ECO - esterified canola oil

Resistance management

Resistance to Sharpen and other MoA Group 14 herbicides could develop through excessive use. Use strictly in accordance with label instructions and in a herbicide programme that incorporates other MoA groups. Refer to the NZCPR website www.resistance.nzpps.org for more detailed information.



For more information on **Sharpen Herbicide**, scan this QR code or contact your local BASF representative on **0800 558 399**.

BASF
We create chemistry

ALWAYS READ AND FOLLOW LABEL DIRECTIONS BEFORE USING ANY PRODUCT IN THIS FACT SHEET.

ACVM Registration numbers: Sharpen: P008391, Arietta: P008825, Frontier-P: P007979. © Copyright BASF 2026 © Registered trademark of BASF.
* Registered trademarks. BASFNZAR0005 0326