

Arietta®

Herbicide



Broad-spectrum post-emergence weed control

Arietta®'s effectiveness, flexible application timing and excellent tank-mixing compatibility have made it a standard inclusion in maize and sweetcorn spray programmes.

Q Why should I use Arietta?

A To take control and provide protection against yield-sapping weeds before row closure. Arietta is particularly effective against hard-to-control weeds like summer grasses and velvetleaf.

Q What are the recommended application rates?

A Arietta 200 mL/ha + atrazine 1 L/ha + ECO** 500 mL/100 L of water

Q What oil/adjuvant should I use?

A An ECO is recommended as they provide Arietta with the most effective penetrant system, e.g. Hasten™, Kwickin™ or Bonza® Gold

Q What are some of the tank-mixing options for maize to extend weed control?

A Atrazine - Enhances Arietta performance on labelled weeds

Nicosulfuron (WG formulations only) - Ryegrass and couch (twitch)

Clopyralid - Californian thistle, yarrow and volunteer potatoes

Dicamba - Californian thistle, field bindweed (*Convolvulus arvensis*) and cornbind (*Polygonum convolvulus*)

PLEASE NOTE: Do not use dicamba, clopyralid or nicosulfuron mixtures on sweetcorn

Q What is the recommended water rate?

A Minimum of 200 L/ha. "Coverage is Key" – for the best results use 300-400 L/ha

Q What are the plant back periods for Arietta?

A The plant backs for Arietta are as follows:

Crop	Plantback times
Wheat, Ryegrass, Carrot, Squash, Onions, Tomatoes	2 weeks
Radish	4 weeks
White Clover & peas	8 weeks
Other crops	Pot test prior to planting or seek advice

* ai - active ingredient, ** ECO - esterified canola oil

BASF
We create chemistry

Arietta®

Herbicide

Q Rainfastness?

A Do not apply if rain is expected within 2 hours of application

Q What are the withholding periods for sweetcorn and maize?

A DO NOT apply after the crop has passed the 6-leaf stage (GS16)


Q What is the withholding period for maize silage/green feed?

A Allow 56 days between application and cutting or introduction of stock

Post-emergent broadleaf and grass weed control in maize and sweetcorn

Broadleaf weeds (up to 8 leaves)			Grass weeds (up to 3 tillers)
Apple of Peru	Fathen	Velvetleaf (before it lignifies)	Barnyard grass
Black nightshade	Fleabane	White clover	Smooth witchgrass
Broad-leaved dock (seedling)	Groundsel	Willow weed	Summer grass
Chamomile	Large flowered mallow		Yellow bristle grass
Chickweed	Scrambling speedwell		
Dandelion	Twin cress		

Arietta – recommended application timing



Scan the QR code to find out more about **Arietta Herbicide** or call **0800 558 399**.

ALWAYS READ AND FOLLOW LABEL DIRECTIONS.

This leaflet is intended as general advice. Arietta ACVM Reg. No. P008825, Sharpen ACVM Reg. No. P008391. Hasten™ is a Trademark of the Victorian Chemical Company. Bonza Gold® is a registered trademark of Nufarm, registered pursuant to the ACVM Act 1997, No. P8871 © Copyright BASF 2026 ® Registered trademark of BASF * Registered trademark. BASFNZAR0022 0926.

BASF
We create chemistry